arXiv:1210.1569v1 [physics.gen-ph] 4 Oct 2012

Observers and Velocity Measurements in Static Spherically Symmetric Space-time: A General Relativistic Prescription

Sumanta Chakraborty *
Department Of Physics, Rajabajar Science College,
Calcutta University, 92 A.P.C Bose road, Kolkata-700009, India

The general misconception regarding velocity measurements of test particle as it approaches black hole is addressed by introducing generalised observer set. We also generalise the motion from radial geodesic to motion confined in a specific θ plane exploiting the symmetry of the problem. For a general static spherically symmetric metric we find that velocity of the test particle do not approach that of light at event horizon. However if the metric co-efficient on the 2-sphere is a function of radial co-ordinte, then the motion is changed drastically and we have shown that this has connection with penrose process.

Keywords: Velocity in General Relativity; Observer; Dilaton Gravity; Penrose Process;

I. INTRODUCTION

The radial motion of a test particle falling in a black hole is one of the key issues in general relativity. The infalling motion has been studied specifically for Schwarzchild black hole by several authors (see [1], [2], [3] and [4]). All of them reached the same conclusion that velocity of the infalling particle approaches that of light near the event horizon, which for the Schwarzchild case is at $r = 2M$, where M is the mass of the black hole. The observers, called static observers, are at rest with respect to the mass creating the gravitational field. They are actually the world lines on the hypersurface of orthogonal killing vector field for the metric describing the gravitational field. However there exists a common misconception that particle approaches the speed of light as it moves to the black hole horizon for all observers, but not as a limiting procedure for a static observer at r as $r \to 2m$. However if we assume that the particle approaches the event horizon at the speed of light for a static observer, as we have defined it earlier, then simple velocity composition law tells that it should approach the speed of light for all local observers as space time is locally Minkowskian.

So we have to modify our notion of velocity for a test particle near a black hole for a static observer. This work is done for Schwarzchild black hole by Crawford et. al. (see [5], [6]). The notion of observer is implimented and used in various co-ordinate frames by Bol$o^{'}$s (see [7], [8], [9]).

However recently Chakraborty et. al. (see [10]) has obtained trajectory around a general spherically symmetric non-rotaing black hole by choosing a general metric ansatz,

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^d\Omega^2 \tag{1}$$

For this general case we find the velocity of the test particle with respect to a static observer (r =constnat) to be a function of $f(r)$. While for the case of a general observer such that both the observer and the test particle moves along geodesic in $\theta = \frac{\pi}{2}$ plane then the velocity of the test particle with respect to the observer to our surprise, do not depend on the choice of the function $f(r)$ however it depends on the angular momenta which was absent in the work by Crawford et. al., who assumed only radial geodesics. By including the angular momenta a rich structure will arise as we have shown later. Also we have used a second kind of spherically symmetric solution inspired by dilaton gravity (see [11]) to show that the velocity depends on the functional form of the metric co-efficient before 2-sphere line element. Here we introduce a function of r as the metric co-efficient on the 2-sphere. The metric is thus taken to be,

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + g(r)d\Omega^2 \tag{2}$$

---

* sumantac.physics@gmail.com

We shall show that the zeros of $g(r)$ are related to penrose process. Throughout the paper we shall use natural unit such that $G = c = 1$.

This paper is organized as follows, in section (II) we introduce the general idea of observer and co-ordinate frames which we shall use through out this work.In section (III) we discuss the motion in spherical symmetric space-time with the metric ansatz given in (1) and then in section (IV) we discuss the motion with the general spherically symmetric model given by equation (2). The paper ends with a short discussion.

II. CO-ORDINATE SYSTEM, REFERANCE FRAMES AND OBSERVERS

The mathematical beauty of general relativity is the freedom of choice of co-ordintes in the description of physical phenomenon. We could choose any co-ordinate system as we wish, this choice might be taken in favour of the symmetry involved in the problem. Also the co-ordinates are not sufficient we need referance frame as well. However the co-ordinate system and referance frames are not independent, for example in one referance frame one set of co-ordinates may be important while it could change in other referance frame. However in literature (see [3]) it is often seen that co-ordinate system and referance frames are used interchangebly. However in our discussion we find the use of "referance frame" and "co-ordinate system" to be distinct. By referance frame we shall mean a set of observers to take measurements, for example the set of all observers moving in a time like geodesic form a referance frame, whereas co-ordinate system refer to numbers specified over the whole space time manifold.

In special relativity an infinite lattice work of sticks and clocks (see [12]) suffice to define a unique referance frame. However in general relativity we cannot have such rigid framework since the space time is Minkowskian only locally, so we replace this rigid system by a fluid (see [4]). In a strictly mathematical sense the set of observers represents a set of future pointing timelike congruence, which is a three parameter family of curves $x^{\mu}(\lambda, y^i)$, where λ is an affine parameter defined over the path, and y^i labels the spatial parts of the curve.

Observer in general theory is very local and it is a material particle parametrized by proper time. An observer field i.e. its velocity field u on the manifold M is stationary provided there exist a smooth function f greater than 0, such that $fu = \xi$ is a killing vector field, so the lie derivative of the metric with respect to the vector field ξ vanishes (i.e. $L_{\xi} g_{\mu\nu} = 0$).

There is a natural way for an u-observer to define the speed of any particle with four velocity t^{μ} as it passes an event $p \in M$, then the observer measure the square of the speed at event p to yield (see [5]),

$$v^2 = \frac{(g_{\mu\nu} + u_{\mu}u_{\nu})t^{\mu}t^{\nu}}{(u_{\alpha}t^{\alpha})^2} \tag{3}$$

Then we have $g_{\mu\nu}t^{\mu}t^{\nu} = -1$ and as well as $u_{\mu}u_{\nu}t^{\mu}t^{\nu} = (u_{\alpha}t^{\alpha})^2$. Thus the above relation can be simplified to yield,

$$v^2 = 1 - \frac{1}{(u^{\mu}t_{\mu})^2} \tag{4}$$

Note that the two velocities u^{μ} and t^{μ} are timelike as observer and the test particle are both time like.

III. MOTION IN A GENERAL SPHERICALLY SYMMETRIC SOLUTION-I

A. Test Particle Geodesic

We shall assume that our test particle is confined to a plane which is generally chosen as $\theta = \pi/2$ for calculational simplicity and as well as we have spherical symmetry so if we discuss the situation for some specified θ plane then it would be the same for all. Thus this no longer represent a radially ingoing particle but a more generalised case where the particle has two variables to specify namely, (r, ϕ). The motion is determined by the Euler equations corresponding to the lagrangian formed as $2L = g_{\mu\nu}\dot{x}_{\mu}\dot{x}_{\nu}$, Which has the following explicit form (using equation (1)),

$$2L = -f(r)\dot{t}^2 + \frac{1}{f(r)}\dot{r}^2 + r^2\dot{\phi}^2 \tag{5}$$

where dot denotes differentiation with repect to proper time of the particle. This equation can be written in terms of the particle proper time and then along the orbit we have $2L = -1$. This finally leads to,

$$d\tau^2 = f(r)dt^2 - \frac{1}{f(r)}dr^2 - r^2 d\phi^2 = f(r)dt^2[1 - v^2] \tag{6}$$

where

$$v^2 = \frac{1}{f(r)^2}\left(\frac{dr}{dt}\right)^2 + \frac{r^2}{f(r)}\left(\frac{d\phi}{dt}\right)^2 \tag{7}$$

This is the velocity of the particle with respect to a ststic observer (r=constant) as illustrated by plugging $u^\mu = (1, 0, 0, 0)$ in equation (3); i.e. the particle moves through a distance $\frac{1}{\sqrt{f}}\sqrt{dr^2 + r^2 f d\phi^2}$ in a proper time given by $\sqrt{f}dt$. Where from now on we shall use simply f for $f(r)$ due to notational simplicity.

Since the lagrangian as given in (5) do not contain t explicitly we have a constant of motion which is nothing but the energy of the particle and it is given by,

$$-\frac{\partial L}{\partial \dot{t}} = f\dot{t} = constant = E \tag{8}$$

This constant of motion actualy originates from the killing vector field $\frac{\partial}{\partial t}$, this can be phrased as, if the 4-velocity of the particle t^a is a geodesic, then we have $\triangledown_t t = 0$. From (6) and (8) we have obtained,

$$v = \sqrt{1 - \frac{f}{E^2}} \tag{9}$$

Also the energy can be determined from the initial value of radius and velocity using (9) such that, $E^2 = \frac{f(R)}{1-v_0^2}$. Where R is the initial radial co-ordinte and v_0 is the initial velocity.

We have another constant of motion in this case which corresponds to the angular momentum of the particle and could be given by,

$$\dot{\phi} = \frac{L}{r^2} \tag{10}$$

Thus finally the velocity in proper frame on the plane $\theta = \frac{\pi}{2}$ is given by,

$$\left(\frac{dr}{d\tau}\right)^2 = \left(\frac{dr}{dt}\right)^2\left(\frac{dt}{d\tau}\right)^2 = E^2 - V^2 \tag{11}$$

where $V^2 = f\left[1 + \frac{r^2}{f^2}E^2\left(\frac{d\phi}{dt}\right)^2\right] = f\left[1 + \frac{L^2}{r^2}\right]$.

Thus the 4-velocity components for the geodesic particle specified by energy and angular momentum is given by,

$$t^\mu = \left(\frac{E}{f}, \sqrt{E^2 - V^2}, \frac{L}{r^2}, 0\right) \tag{12}$$

written in terms of the constants of motion E and L. As a check we can use the identity $t_\mu t^\mu = -1$ and from (12) we see that $u_\mu u^\mu = -\frac{E^2}{f} + \frac{E^2 - V^2}{f} + \frac{L^2}{r^2} = -\frac{V^2}{f} + \frac{L^2}{r^2} = -1$. Hence the four velocity as represented by (12) is correct.

B. Static Limit

In some cases the velocity is measured in terms of proper time, as determined by clocks synchronized along trajectory of the particle. The velocity in case of radial particle is given by (see [1]),

$$v^2 = \left(g_{00} + g_{01}\frac{dx^1}{dx^0}\right)^{-2} \left(g_{01}^2 - g_{00}g_{11}\right) \left(\frac{dx^1}{dx^0}\right)^2 \tag{13}$$

When we generalise this result to our case where we have three co-ordinats x^0, x^1 and x^3 (since $x^2 = \theta = constant$), then velocity expression generalises to,

$$v^2 = \frac{\left(g_{10}^2 - g_{00}g_{11}\right)\left(\frac{dx^1}{dx^0}\right)^2 + \left(g_{30}^2 - g_{00}g_{33}\right)\left(\frac{dx^3}{dx^0}\right)^2 + 2\left(g_{10}g_{30} - g_{13}g_{00}\right)\frac{dx^1}{dx^0}\frac{dx^3}{dx^0}}{\left(g_{00} + g_{10}\frac{dx^1}{dx^0} + g_{30}\frac{dx^3}{dx^0}\right)^2} \tag{14}$$

Note that if we let $\frac{dx^3}{dx^0}$ to be zero, then it reduces to equation (13). In our case kepping the non zero terms we obtain,

$$v^2 = \frac{1}{f^2}\left(\frac{dr}{dt}\right)^2 + \frac{r^2}{f}\left(\frac{d\phi}{dt}\right)^2 \tag{15}$$

which is completely identical to (7). This definition has co-ordinte invariance. The 4 velocity has components $u_\mu = (-g_{00})^{-1/2}g_{\mu 0}$ and that for the particle reduces to $t^\mu = (\frac{dx^0}{d\tau}, \frac{dx^1}{d\tau}, 0, \frac{dx^3}{d\tau})$. Thus using (4) we obtain the same equation as (15).

From (9) we see that as $f(r) = 0$, the velocity is equal to 1. Hence for static observers v appproaches the speed of light at the event horizon and they predict faster than light speed inside event horizon.

It might seem at first sight that this result has nothing to do with $f(r) = 0$ but is connected to the co-ordinate system. However it has nothing to do with co-ordinate system but with the observer. So we should generalise our observer set.

Also no observer can be at rest at $r = 2M$ except photon, with respect to photon all particle traverse at speed of light. To get a clear view we discuss the acceleration of a static observer in the field of the gravitating body. The acceleration is necessary as in general relativity an observer at rest is not geodesic and is accelerated.

The four acceleration field is defined as,

$$a^\eta = u^\eta_{;\mu}u^\mu = \left(u^\eta_{,\mu} + u^\alpha \Gamma^\eta_{\alpha\mu}\right) \tag{16}$$

The only non zero component is given by using the definition of four velocities for static observers , $u_\mu = \frac{g_{\mu 0}}{\sqrt{-g_{00}}}$ to yield,

$$a^1 = \frac{1}{2}\frac{df}{dr} \tag{17}$$

So acceleration depends on the function $f(r)$.

C. Ingoing Observers

We consider motion of two particles such that the four velocities are given by,

$$\left.\begin{array}{l} t^\mu = \left(\frac{E_1}{f}, \sqrt{E_1^2 - V_1^2}, 0, \frac{L_1^2}{r^2}\right) \\ u^\nu = \left(\frac{E_2}{f}, \sqrt{E_2^2 - V_1^2}, 0, \frac{L_2^2}{r^2}\right) \end{array}\right\} \tag{18}$$



Figure 1: The figure shows variation of v^2 with radial co-ordinate r for different choice of E_1, E_2, L_1 and L_2.



Figure 2: The figure shows variation of v^2 with test particle energy for diferent observer energy and radial distance.



Figure 3: The figure shows variation of v^2 with test particle angular momentum for differnt choices of test particle energy



Figure 4: The figure shows variation of v^2 with E_2 and r.

Hence we obtain the following result,

$$t^{\mu}u_{\mu} = g_{\mu\nu}t^{\mu}u^{\nu} = -\frac{E_1E_2}{f} + \frac{\sqrt{\left(E_1^2 - V_1^2\right)\left(E_2^2 - V_1^2\right)}}{f} + \frac{L_1L_2}{r^2} \tag{19}$$

Thus we obtain,

$$\left(t^{\mu}u_{\mu}\right)^2 = \left(\frac{E_1E_2}{f}\right)^2 \left[1 - \frac{fL_1L_2}{r^2E_1E_2} - \sqrt{\left(1 - fa_1\right)\left(1 - fa_2\right)}\right]^2 \tag{20}$$

Where, $a_1 = \frac{1}{E_1^2}\left(1 + \frac{L_1^2}{r^2}\right)$ and $a_2 = \frac{1}{E_2^2}\left(1 + \frac{L_2^2}{r^2}\right)$.

Simplifying and rearranging terms we have obtained that,

$$\left(t^{\mu}u_{\mu}\right)^2 = E_1^2E_2^2\left[\frac{1}{2}\left(a_1 + a_2\right) - \frac{L_1L_2}{r^2E_1E_2}\right]^2 \tag{21}$$

We know that the velocity could be given by, $v^2 = 1 - \frac{1}{\left(u^{\mu}t_{\mu}\right)^2}$. Thus using equation (21) we obtain after simplification,

$$v^2 = 1 - \frac{4}{E_1^2E_2^2\left[\left(\frac{1}{E_1^2} + \frac{1}{E_2^2}\right) + \frac{1}{r^2}\left(\frac{L_1}{E_1} - \frac{L_2}{E_2}\right)^2\right]^2} \tag{22}$$

Note that as $r \to 0$ the velocity approaches that of light i.e. $v = 1$. However if the particle and the observer has the same impact parameter i.e. $\frac{L_1}{E_1} = \frac{L_2}{E_2}$ then even if $r \to 0$ the velocity does not approach zero, which is a very interesting result.

Note from figure-1, as radial co-ordinate of the particle is decreased the veloity remain less than speed of light. As $r \to 0$ the velocity also approaches 1 in our system of units, which is justified and shows the actual motion that happen as the particle moves within the event horizon. It is also clear that with increase of the energy of the particle the velocity increases and it also increases with increasing the angular momenta.

From figure-2 we see that as energy of the particle is increased we get a interesting behaviour, at first it decreases and become zero, then it again increases. Thus here the combined quantity in the denominator becomes 4. This happens when E_1 coincides with E_2 (see the figure), as we have chosen $L_1 = L_2$ (see equation 22). However changing the radius has a very small effect on velocity profile.

From figure-3 we find that velocity varies with angular momentum in some what the same manner as it does with energy. However by proper choice of $E_2 = E_1$ the velocity can be made zero when $L_2 = E_2$ as we have chosen other parameters such that $L_1 = E_1$, since under this condition the denominator in equation (22) become 4. As well as we can eliminate that zero by changing E_2.

Figure-4 shows the variation of velocity both with radial co-ordinate and the energy of the particle. This graph merely shows combined effects of varying radius and energy as we have illustrated in earlier graphs.

IV. MOTION IN A GENERAL SPHERICALLY SYMMETRIC SOLUTION-II

In this section we shall consider general spherically symmetric metric (given by equation 2) inspired by dilaton gravity. Thus our lagrangian reads,

$$2L = -f\dot{t}^2 + \frac{1}{f}\dot{r}^2 + g\dot{\phi}^2 \tag{23}$$

As usual we have $f = f(r)$ and $g = g(r)$, however due to notational simplicity we have taken them to be simply f and g respectively. Then the velocity has the following expression,

$$v^2 = \frac{1}{f^2}\left(\frac{dr}{dt}\right)^2 + \frac{g}{f}\left(\frac{d\phi}{dt}\right)^2 \tag{24}$$

The potential has the following expression which could be given by,

$$V^2 = f\left(1 + \frac{L^2}{g}\right) \tag{25}$$

Thus 4-velocity components are given by for this potential to yield,

$$u^\mu = \left(\frac{E}{f}, -\sqrt{E^2 - V^2}, 0, \frac{L}{g}\right) \tag{26}$$

Note that for this case as well we have the following result $t^\mu t_\mu = -1$. If we have used equation (14) then we might have obtained that the velocity has the same expression as that given by (24).

Also the acceleration has no change only a^1 is non zero and has the value given by (17). If we proceed in an identical way then we obtain the following result for the velocity of a particle relative to an observer,

$$v^2 = 1 - \frac{4}{E_1^2 E_2^2\left[\left(\frac{1}{E_1^2} + \frac{1}{E_2^2}\right) + \frac{1}{g}\left(\frac{L_1}{E_1} - \frac{L_2}{E_2}\right)^2\right]^2} \tag{27}$$

The most interesting part of this velocity expression corresponds to the fact that when at some finite r the quantity $g = 0$ then $v = 1$. So even if it had not go to $r = 0$ the particle is seen to move with the velocity of light. However under the same situation as above such that both the particle and the observer moves with the same impact parameter we obtain that this case is prohibited and the particle has v less than 1 for all r.

Now to illutrate we shall use a specific form for $g(r)$ from dilaton gravity. To give a brief introduction to dilaton grvity we point out some basic notion regarding this black hole solution. Static uncharged black hole in general relativity are described by Schwarzschild solution. If mass of the black hole is much large compared to Planck mass then this also describes to a good approximation the uncharged black hole in string theory except regions near singularity. However this is not the case for Einstein-Maxwell solutions. The dilaton coupling with F^2 implies that every solution with non zero $F_{\mu\nu}$ will come with a non zero dilaton. Thus the charged black hole solution in general relativity (which is the Reissner-nordström solution) appears in a new form in string theory due to the presence of dilaton. The effective four dimensional low energy lagrangian obtained from string theory is,

$$S = \int d^4x\sqrt{-g}[-R + e^{-2\phi}F^2 + 2(\nabla\phi)^2]$$

where $F_{\mu\nu}$ is the Maxwell field associated with a $U(1)$ subgroup of $E_8 \times E_8$ or $Spin(32)/Z_2$. We have set the remaining gauge fields and antisymmetric tensor field $H_{\mu\nu\rho}$ to zero and ϕ represents the dilaton strength (see Garfinkle et.al.[11], Coleman [13],, Bekenstein [14] and Witten [15])

The static spherically symmetric solution would give the following line element corresponding to the above action as,(see Garfinkle et.al.[11])

$$ds^2 = -(1 - \frac{2M}{r})dt^2 + \frac{1}{(1 - \frac{2M}{r})}dr^2 + r(r - e^{2\phi_0}\frac{Q^2}{M})d\Omega^2 \tag{28}$$

Where ϕ_0 is the asymptotic value of dilaton strength. Thus applying the value of $g(r)$ from (28) we could obtain the velocity expression from (27). We shall now present our results for this choice graphically.

From figure-5 we see that velocity of the particle has similar kind of variation with energy of the particle as we have seen in previous section. However note that even if we change the dilaton strength the zero



Figure 5: The figure shows variation of v^2 with E_2 for differnt choices of dilaton strength f_0.



Figure 6: The figure shows variation of v^2 with L_2 for different choices of test particle energy E_2 and dilaton strength f_0.



Figure 7: The figure shows variation of v^2 with L_2 for different choices of dilaton Strength f_0 for a specified radius such that $g(r)$ is greater than zero always.



Figure 8: The figure shows variation of v^2 with r for different choices of test particle energy E_2 and angular momentum L_2.



Figure 9: The figure shows variation of v^2 with L_2 and E_2, the test particle energy and angular momentum.



Figure 10: The figure shows variation of v^2 with r and dilaton strength f_0.

in the velocity function do not shift. Figure-6 shows a very interesting feature. At large value of angular momentum the velocity approaches that of 1 in our system of units, which is justified. However note that a given region of space has no velocity which mean that we can not probe that region, with increasing dilaton strength the forbidden region increases. This is the sign that we have a high energy barrier. This comes from the zero of the function $g(r)$. This is also a horizon except the event horizon. This is what influence the velocity of the test particle. So note that the region within $r = 2M$ and $r = e^{2\phi_0}\frac{Q^2}{M}$ is responsible for this behaviour. Here both the particle and the observer took energy from the black hole so that they no longer follow the geodesic path, so that region is singular region of our velocity expression. However within $r = 2M$ the particle and observer behave properly, giving correct velocity expression. This phenomenon is similar to what we see for penrose process. Thus this onset of forbidden region actually shows the penrose process in action. However if we tune our conditions such that the quantity $e^{2\phi_0}\frac{Q^2}{M}$ is less than the radial co-ordinate we have no such discontinuity as the particle and the observer have not reached the radius where $g(r) = 0$ (see figure-7). Then the variation of velocity with radius is shown in figure-8, figure-9 shows variation of velocity with both energy and angular momentum of the test particle. Figure-10 shows the variation of velocity with radius and dilaton strength, which also show spikes at the position of (r, ϕ_0) such that the relation $r = e^{2\phi_0}\frac{Q^2}{M}$ is satisfied. Which again shows the penrose process giving energy to the particle and making its velocity 1, i.e. the speed of light.

V. DISCUSSION

We have seen that velocity of any ingoing particle with respect to observer sets as defined in the section (II) for a general spherically symetric potential with unit 2-sphere is always less than that of light outside the singular point, it approaches the speed of light as $r \rightarrow 0$. However the notion of static observers are not valid for $r \leq 2M$. It is valid only for region outside the event horizon. Thus we have defined ingoing observers and determine veloity with respect to the observer. We found that velocity of the test particle always remain less than 1. For a different choice of metric with a function on 2-sphere we found that the velocity is always less than 1 but there are some region which is forbidden. This represents the onset of penrose process. The region where this process becomes important corresponds to $2M \leq r \leq e^{2\phi_0}\frac{Q^2}{M}$. So from determining the velocity of a test particle we could understand many diverse processes like penrose process, which we have just elaborated. Also it should be noted that the above analysis is not restricted to Einstein gravity, it can be applied to other black hole solutions in modified gravity theories. Also it could be extended to higher dimension.

Acknowledgement: The author thanks prof. Subenoy Chakraborty of Jadavpur University for helpfull discussion. The author also thanks DST, Govt. of India for awarding KVPY fellowship. He thanks IUCAA, Pune, for warm hospitality where a part of this work was done.

[1] Landau, L. and Lifschitz, E.:- (1971) **The Classical Theory of Fields**, 3rd ed. (*Addison-Wesley, Reading, Massachusetts*).

[2] Wald, R. M.:- (1984) **General Relativity** (*University of Chicago Press, Chicago, Illinois*).

[3] Bergmann, P. G.:- (1942) **Introduction to The Theory of Relativity** (*Prentice-Hall, New York*).

[4] Moller, C.:- (1972) **The Theory of Relativity**, 2nd ed. (*Oxford University Press, Delhi*).

[5] Crawford, P. and Tereno, I.:- *Gen. Relativity Gravitation* **34** (2002), 2075.

[6] Janis, A.:- *Phy. Rev. D* **15** (1977), 3068.

[7] Bolo′s, V.J. *preprint* **gr-qc/0506032v4** (2011).

[8] Ellis, G. F. R., Nel, S. D., Maartens, R., Stoeger, W. R. and Whitman, A.P.:- *Phys. Rep.* **124** (1985), 315.

[9] Bol$o^{'}$s, V. J.:- *J. Geom. Phys.* **56** (2006), 813.

[10] Chakraborty, Sumanta. and Chakraborty Subenoy.:- *Can. J. Phys.* **89** (2011), 689 [arXiv:1109.0676[gr-qc],2011].

[11] Garfinkle, D., Horowitz, G. T. and Strominger, A.:- Phys. Rev. D **43**, 3140 (1991).

[12] Teller, E.F. and Wheeler, J. A.:- (1992) **Spacetime Physics: Introduction to Special Relativity** (*W. H. Freeman, San Fransisco*).

[13] Coleman, S.:- (1983) in *The Unity of the Fundamental Interactions*, edited by A.Zichichi (Plenum, London).

[14] Bekenstein, J.:- Phys. Rev. D **5**, 1239 (1972); Ann.phys. (N.Y.) **91**, 75 (1975);Bocharova, N.,Broonikov, K. and Melnikov, V. Vestn.Mosk.Univ.Fiz.Astron. **6**, 706 (1970).

[15] Witten (ed.) (1962) *Gravitation: An Introduction to Current Reaserch*, (Wiley, N.Y.).